*Grass and Forage Science* (1989) 44, 451–454

# Author index

AKIN D.E. *see* WILSON J.R.
ANDERSON R. *see* KENNEDY S.J.
ANDERSON R. *see* STEEN R.W.J.
ANDERSON R., GRACEY H.I., KENNEDY S.J., UNSWORTH E.F. and STEEN R.W.J. Evaluation studies in the development of a commercial bacterial inoculant as an additive for grass silage. 1. Using pilot-scale silos, 361
ARMSTRONG R.H., COMMON T.G. and DAVIES G.J. The prediction of the *in vivo* digestibility of a diet of sheep and cattle grazing indigenous hill plant communities by *in vitro* digestion, faecal nitrogen concentration or 'indigestible' acid-detergent fibre, 303
BAKER R.D. *see* GIBB M.J.
BARNARD D.J. BOOK REVIEW. KELLY A.F. Seed Production of Agricultural Crops, 359
BARTHRAM G.T. *see* GRANT SHEILA A.
BELL, C.C. and RITCHIE I.M. The effect of frequency and height of defoliation on the production and persistence of 'Grasslands Matua' prairie grass (Research Note), 245
BENNET D.W. *see* GUPTA M.L.
BERGSMA K. *see* VAN VUUREN A.M.
BLACKBURN H.D. and KOTHMANN M.M. A forage dynamics model for use in range or pasture environments, 283
BOUCAUD J. *see* GONZALEZ B.
BRADLEY N.W. *see* DOUGHERTY C.T.
BROOKFIELD S.K.E. *see* TALLOWIN J.R.B.
BROOKMAN S.K.E *see* TALLOWIN J.R.B.
CABALLERO R. Yields and chemical composition of whole-crop field beans and their components during pod-filling (Research Note), 347
CEBRIAN MERCEDES *see* OSORO K.
COLLINS ROSEMARY P. & RHODES I. Yield of white clover populations in mixture with contrasting perennial ryegrasses (Research Note), 11
COMMON T.G *see* ARMSTRONG R.H.
CORBETT J.L. *see* WHEELER J.L.
CORNELIUS P.L. *see* DOUGHERTY C.T.
DAVIES D.A., FOTHERGILL M. and JONES D. Assessment of contrasting perennial ryegrasses, with and without white clover, under continuous sheep stocking in the uplands. 1. Animal production from the grass varieties, 431
DAVIES D.A., FOTHERGILL M. and JONES D. Assessment of contrasting perennial ryegrasses, with and without white clover, under continuous sheep stocking in the uplands. 2. The value of white clover for lamb production, 441
DAVIES D.A., FOTHERGILL M. and JONES D. Frequency of stocking rate adjustments required on contrasting upland perennial ryegrass pastures continuously grazed to a sward height criterion from May to July, 213
DAVIES G.J. *see* ARMSTRONG R.H.
DOUGHERTY C.T., BRADLEY N.W., CORNELIUS P.L. and LAURIAULT L.M. Ingestive behaviour of beef cattle offered different forms of lucerne (*Medicago sativa* L.), 335
DOUGHERTY C.T., BRADLEY N.W., CORNELIUS P.L. and LAURIAULT L.M. Short-term fasts and the ingestive behaviour of grazing cattle, 295
DUYME M. *see* GONZALEZ B.
ELSTON D.A. *see* GRANT SHEILA A.
ESSLEMONT R.E. *see* WILTSHIRE J.J.J.
ESTELL R.E. *see* HOLECHEK J.L.
FENLON D.R., WILSON J. and WEDDELL D.R. The relationship between spoilage and *Listeria monocytogenes* in bagged and wrapped big bale silage (Research Note), 97
FOTHERGILL M. *see* DAVIES D.A.
FRAME J. Herbage productivity of a range of grass species under a silage cutting regime with a high fertilizer nitrogen application, 267
FRAME J. *see* KLESSA D.A.
FROL-KRAMER F. *see* VAN VUUREN A.M.
GALYEAN M.L. *see* HOLECHEK J.L.

GIBB M.J. and BAKER R.D. Effect of changing grazing severity on the composition of perennial ryegrass/white clover swards stocked with beef cattle, 329
GIBB M.J., BAKER R.D. and SAYER ANNA M.E. The impact of grazing severity on perennial ryegrass/white clover swards stocked continuously with beef cattle, 315
GOLIGHTLY R.D. *see* KLESSA D.A.
GONZALEZ B., BOUCAUD J., SALETTE J., LANGLOIS J. and DUYME M. Changes in stubble carbohydrate content during regrowth of defoliated perennial ryegrass (*Lolium perenne* L.) on two nitrogen levels, 411
GORDON F.J. A further study on the evaluation through lactating cattle of a bacterial inoculant as an additive for grass silage (Research Note), 353
GORDON F.J. An evaluation through lactating cattle of a bacterial inoculant as an additive for grass silage, 169
GRACEY H.I. *see* ANDERSON R.
GRACEY H.I. *see* KENNEDY S.J.
GRACEY H.I. *see* STEEN R.W.J.
GRANT SHEILA A., ELSTON D.A. and BARTHRAM G.T. Problems of estimating tissue turnover in grass swards in the presence of grazing animals, 47
GUPTA M.L., MACMILLAN R.H., MCMAHON T.A. and BENNETT D.W. A simulation model to predict the drying time for pasture hay, 1
HAGGAR R.J., ISAAC S.P. and STANDELL C.J. Modifying growth patterns in nine grasses with mefluidide (Research Note), 85
HARKESS R.D. *see* KLESSA D.A.
HAY R.K.M. The performance of high-latitude grass varieties under Scottish conditions: seasonal distribution of dry-matter production, 405
HEBBLETHWAITE P.D. *see* WILTSHIRE J.J.J.
HIDES D.H. *see* HOLLINGTON P.A.
HIDES D.H. *see* MARSHALL A.H.
HILL M.J. Growth of *Trifolium repens* L. and *Trifolium semipilosum* Fres. var. *glabrescens* Gillet at different temperatures in controlled environments and in the field, 125
HOLECHEK J.L., ESTELL R.E., GALYEAN M.L. and RICHARDS W. Chemical composition, *in vitro* digestibility and *in vitro* VFA concentrations of New Mexico native forages (Research Note), 101
HOLLINGTON P.A. *see* MARSHALL A.H.
HOLLINGTON P.A., MARSHALL, A.H. and HIDES D.H. Effect of seed crop management on potential seed yield of contrasting white clover varieties. II. Seed yield components and potential seed yield, 189
HOPKINS A., MURRAY P.J. and PATEFIELD W.M. A comparison of the herbage productivity of *Bromus wildenowii* cv. Grasslands Matua with four cultivars of *Lolium perenne* when grown in association with *Trifolium repens*, 31
HUMPHREYS M.O. Water-soluble carbohydrates in perennial ryegrass breeding. I. Genetic differences among cultivars and hybrid progeny grown as spaced plants, 231
HUMPHREYS M.O. Water-soluble carbohydrates in perennial ryegrass breeding. II. Cultivar and hybrid progeny performance in cut plots, 237
HUMPHREYS M.O. Water-soluble carbohydrates in perennial ryegrass breeding. III. Relationships with herbage production, digestibility and crude protein content, 423
ISAAC S.P. *see* HAGGAR R.J.
JONES D *see* DAVIES D.A.
JONES E.L. and ROBERTS J.E. Sward maintenance of *Lolium multiforum* by slurry seeding, 27
KEILLER P.R. *see* THOMPSON N.
KENNEDY S.J. *see* ANDERSON R.
KENNEDY S.J. *see* STEEN R.W.J.
KENNEDY S.J., GRACEY H.I., UNSWORTH E.F., STEEN R.W.J. and ANDERSON R. Evaluation studies in the development of a commercial bacterial inoculant as an additive for grass silage. 2. Responses in finishing cattle, 371
KILPATRICK D.J. *see* STEEN R.W.J.
KLESSA D.A., FRAME J., GOLIGHTLY R.D. and HARKESS R.D. The effect of fertilizer sulphur on grass production for silage, 277
KOTHMANN M.M. *see* BLACKBURN H.D.
LAIDLAW A.S. and STEEN R.W.J. Turnover of grass laminae and white clover leaves in mixed swards continuously grazed with steers at a high- and low-N fertilizer level, 249
LANGLOIS J. *see* GONZALEZ B.
LAURIAULT L.M. *see* DOUGHERTY C.T.
LEWIS G.C. Factors influencing the effects of fungicide treatement on seedling emergence of perennial ryegrass (*Lolium perenne* L.), 417
MACMILLAN R.H. *see* GUPTA M.L.
MARSHALL A.H. *see* HOLLINGTON P.A.

MARSHALL A.H., HOLLINGTON P.A. and HIDES D.H. Effect of seed crop management on the potential seed yield of contrasting white clover varieties. I. Influorescence production, 181
MCGILLOWAY D.A. *see* WILTSHIRE J.J.J.
MCLEOD M.N. *see* WILSON J.R.
MCMAHON T.A. *see* GUPTA M.L.
MINSON D.J. *see* WILSON J.R.
MOSELEY G. and RAMANATHAN V. The effect of dry feed additives on the nutritive value of silage, 391
MOWAT D.J. and SHAKEEL M.A. The effect of different cultivars of clover on numbers of, and leaf damage by, some invertebrate species, 11
MOWAT D.J. and SHAKEEL M.A. The effect of some invertebrate species on persistence of white clover in ryegrass swards, 177
MUCK R.E. Initial bacterial numbers on lucerne prior to ensiling, 19
MURRAY P.J. *see* HOPKINS A.
OSORO K. and CEBRIAN MERCEDES. Digestibility of energy and gross energy intake in fresh pastures, 41
PATEFIELD M. *see* TALLOWIN J.R.B.
PATEFIELD W.M. *see* HOPKINS A.
RAMANATHAN V. *see* MOSELEY G.
RANGELEY ANNE. The suitability of plant tissue tests to indicate phosphorus deficiency in perennial ryegrass (*Lolium perenne* L.) (Research Note), 91
RHODES I. *see* COLLINS ROSEMARY P.
RICHARDS W. *see* HOLECHEK J.L.
RITCHIE I.M. *see* BELL C.C.
ROBERTS D.J. A comparison of buffer and partial storage feeding of a straw/concentrate mixture to grazing dairy cows, 399
ROBERTS J.E. *see* JONES E.L.
SALETTE J. *see* GONZALEZ B.
SANDERSON M.A. and WEDIN W.F. Nitrogen concentrations in the cell wall and lignocellulose of smooth bromegrass herbage, 151
SANDERSON M.A. and WEDIN W.F. Nitrogen in the detergent fibre fractions of temperature legumes and grasses, 159
SAYER ANNA M.E. *see* GIBB M.J.
SHAKEEL M.A. *see* MOWAT D.J.
STANDELL C.J. *see* HAGGAR R.J.
STEEN R.W.J. *see* ANDERSON R.
STEEN R.W.J. *see* KENNEDY S.J.
STEEN R.W.J. *see* LAIDLAW A.S.
STEEN R.W.J., UNSWORTH E.F., GRACEY H.I., KENNEDY S.J., ANDERSON R. & KILPATRICK D.J. Evaluation studies in the development of a commercial bacterial inoculant as an additive for grass silage. 3. Responses in growing cattle and interaction with protein supplementation, 381
TALLOWIN J.R.B. and BROOKMAN S.K.E. The use of irradiation for the preservation of fresh herbage for subsequent botanical and chemical analysis (Research Note), 343.
TALLOWIN J.R.B., TCACENCO F., PATEFIELD M. and BROOKFIELD S.K.E. A correction factor for the influence of changes in lamina weight per unit length in grasses on measurements of the weight of lamina removed by grazing, 205
TCACENCO F. *see* TALLOWIN J.R.B.
THOMPSON N., KEILLER P.R. and YATES C.W. Predicting the digestibility of grass grown for first-cut silage, 195
UNSWORTH E.F. *see* ANDERSON R.
UNSWORTH E.F. *see* KENNEDY S.J.
UNSWORTH E.F. *see* STEEN R.W.J.
VAN BEERS J.A.C. *see* VAN VUUREN A.M.
VAN VUUREN A.M., BERGSMA K., FROL-KRAMER F. and VAN BEERS J.A.C. Effects of addition of cell wall degrading enzymes on the chemical composition and the *in sacco* degradation of grass silage, 223
VICKERY P.J. *see* WHEELER J.L.
WEDDELL D.R. *see* FENLON D.R.
WEDIN W.F. *see* SANDERSON M.A.
WHEELER J.L. and CORBETT J.L. Criteria for breeding forages of improved feeding value: results of a Delphi survey, 77
WHEELER J.L. and VICKERY P.J. Variation in HCN potential among cultivars of white clover (Trifolium repens) (Research Note), 107
WHYTE T.K. *see* WRIGHT I.A.
WILSON J. *see* FENLON D.R.
WILSON J.R., AKIN D.E., MCLEOD M.N. and MINSON D.J. Particle size reduction of the leaves of a tropical and a temperate grass by cattle. II. Relation of anatomical structure to the process of leaf breakdown through chewing and digestion, 65
WILSON J.R., MCLEOD M.N. and MINSON D.J. Particle size reduction of the leaves of a tropical and a temperate grass by cattle. I. Effect of chewing during eating and varying times of digestion, 55

WILTSHIRE J.J.J., HEBBLETHWAITE P.D., ESSLEMONT R.E. and MCGILLOWAY D.A. The use of growth regular RSWO411 in seed crops of *Lolium perenne* L., 139
WRIGHT I.A. and WHYTE T.K. Effects of sward surface height on the performance of continuously stocked spring-calving beef cows and their calves, 259
YATES C.W. *see* THOMPSON N.